arXiv:1207.7108v2 [math.PR] 19 Aug 2012

Horton self-similarity of Kingman's coalescent tree

Yevgeniy Kovchegov*and Ilya Zaliapin†

Abstract

The paper establishes Horton self-similarity for tree representation of Kingman's coalescent process and, equivalently, level-set tree of a white noise. The proof is based on a Smoluchowski-type system of ordinary differential equations for the number of Horton-Strahler branches in a tree that represents Kingman's coalescent. This approach allows us to establish a weak form of the Horton law, namely the existence of a limit of $(N_k/N)^{1/k}$ for the number N_k of branches of Horton-Strahler order k in a coalescent system of size N, as we let $N \to \infty$ and then $k \to \infty$. We conjecture, based on numerical observations, that the Kingman's coalescent is also Horton self-similar in regular strong sense with Horton exponent $R = \lim\limits_{k\to\infty}\left(\lim\limits_{N\to\infty}(N_{k+1}/N_k)\right) = 0.328533...$ and asymptotically Tokunaga self-similar. Finally, we demonstrate combinatorial equivalence between the trees of a Kingman's coalescent and level-set trees of a discrete white noise.

1 Introduction

When studying the tree graphs associated with random structures one often aims at discovering a particular symmetry or a consistent pattern such as self-similarity. There exist two important types of tree self-similarity related to the Horton-Strahler [11, 39] and Tokunaga [41] indexing schemes for tree branches (see Sect. 4 for definitions). Introduced in hydrology in the 1960s to describe the dendritic structure of river networks, which is among the most evident examples of natural branching, these indexing schemes have been rediscovered and used in other applied fields. The Horton-Strahler indexing assigns orders to the tree branches according to their relative importance in the hierarchy. For instance, in a perfect binary tree of depth $d_{\max}$ the Horton-Strahler order r is uniquely determined by the node depth d: $r = d_{\max} - d + 1$. Hence, all leaves have order 1 and the root has the maximal order $d_{\max} + 1$. The Horton-Strahler orders have been shown useful in diverse applications that range from ranking river tributaries to analysis of brain structure to designing optimal computer codes; see [11, 39, 14, 47, 29, 5, 24, 43, 4, 44, 6] and references therein. Quite often, observed

---

*Department of Mathematics, Oregon State University, Corvallis, OR 97331-4605, USA kovchegy@math.oregonstate.edu

†Department of Mathematics and Statistics, University of Nevada, Reno, NV, 89557-0084, USA zal@unr.edu

systems exhibit geometric decrease of the numbers N_k of elements of Horton-Strahler order $k \geq 1$; this property is called *Horton self-similarity.* The common ratio R of the respective geometric series is called the *Horton exponent.* In accordance with Horton-Strahler rules (see Sect. 4.2, Eq.(2)), a branch of order $k > 1$ can only be created by merging of two branches of orders $(k-1)$. The Horton self-similarity hence addresses the *principal branching* in a tree and presents a counterpart of the power-law element size distribution. The last statement becomes clear if one naturally interpret the Horton-Strahler order k as a logarithmic measure of element size s: $k \approx \log s$. A stronger *Tokunaga self-similarity* addresses so-called *side branching* – merging of branches of distinct orders. The Tokunaga self-similarity implies that different levels of a hierarchical system have the same statistical structure, in a sense to be defined; it is parameterized by a positive pair (a, c).

A classical model that exhibits Horton and Tokunaga self-similarity is critical binary Galton-Watson tree [29, 4, 31], also known in hydrology as Shreve's random topology model [36, 37]; it is conditionally equivalent to the uniform distribution on planar binary trees with N leaves [31]. This model has $R = 4$ and $(a, c) = (1, 2)$. For long time, the critical binary Galton-Watson tree has remained the only well-known probability model for which the Horton and Tokunaga self-similarity was rigorously established, and whose Horton-Strahler ordering has received attention in the literature [36, 37, 14, 47, 5, 29, 27, 4]. The model has been particularly popular in hydrology as an approximation to the topology of the observed river networks [40]. Scott Peckham [29] has first explicitly noticed, by performing a high-precision extraction of river channels for Kentucky River, Kentucky and Powder River, Wyoming, that the Horton exponents and Tokunaga parameters for the observed rivers significantly deviate from that for the Galton-Watson model. He reported values $R \approx 4.6$ and $(a, c) \approx (1.2, 2.5)$ and emphasized the importance of studying broad range of Horton exponents and Tokunaga parameters. The general interest to fractals and self-similar structures in natural sciences during the 1990s led to a quest, mainly inspired and led by Donald Turcotte, for Tokunaga self-similar tree graphs of diverse origin. As a result, the Horton and Tokunaga self-similarity, with a broad range of respective parameters, have been empirically or rigorously established in numerous observed and modeled systems, well beyond river networks. This includes vein structure of botanical leaves [24, 43], diffusion limited aggregation [26, 21], two dimensional site percolation [42, 46, 50, 51], nearest-neighbor clustering in Euclidean spaces [45], a general hierarchical coagulation model of Gabrielov *et al.* [9] introduced in the framework of self-organized criticality, and a random self-similar network model of Veitzer and Gupta [44] developed as an alternative to the Shreve's random topology model for river networks. The increasing empirical evidence prompted the question: *What basic probability models can generate Tokunaga self-similar trees with a range of parameters?* An illuminating result in this direction has been obtained by Burd et al. [4]; these authors have demonstrated that Tokunaga self-similarity is a characteristic property of the critical binary branching in the broad class of (not necessarily binary) Galton-Watson processes. Hence, generic branching cannot explain the variability of Horton and Tokunaga parameters seen in nature. Recently, the authors of the current paper established Horton and Tokunaga self-similarity for the level-set tree representation of a homogeneous discrete Markov chain and infinite-tree representation of a regular Brownian motion in continuous time [49]. The results in [49] (i) expand the class of Horton and Tokunaga self-similar processes beyond the critical binary Galton-Watson branching, since the tree representation of Markov chains in general is not equivalent to the Galton-Watson process [31, 49], and (ii) suggest that coalescent (aggregation) phenomena may be better suited for generating self-similar trees. This study is a first step towards exploring self-similarity of trees generated by coalescent processes.

In general, inferring the dynamics, coalescence (aggregation) versus headward growth (branching), for a static network pattern is non-trivial and largely unsettled; see [12, 18, 10] for both models and empirical approaches. In this regard, it is of interest to analyze the behavior of self-similarity statistics, such as Horton ratios, for both classes of models. The results in this paper focus on coalescence and are quite distinct from those covered by the bulk of the mathematical theory to date.

Coalescent processes have been much studied in diverse scientific areas as well as in applied probability literature. We refer to David Aldous [2] for a comprehensive review and further references, and only mention here that coalescents offer an important tool for studying inheritance relationships in population genetics [15, 16, 17, 30], formation of planets and galaxies in astronomy [32, 33], fracturing of solids [3, 34], development of drainage basins, including extraterrestrial ones [35, 13], dynamics of aerosols [7], kinetics of polymerization and gelation [52], algae growth [1], evolutionary driven aggregation of animals [28] and many other phenomena. Generally speaking, a coalescent describes the process of merging of particles into consecutively larger clusters. If the coalescence rate $K(x, y)$ of two clusters depends only on their masses (sizes) x and y, then the mean-field dynamics of a coalescent process is described by the celebrated Smoluchowski integro-differential equation [38, 2] (see also Sect. 3). The most studied in this class are the Kingman's ($K(x, y) = 1$), additive ($K(x, y) = x + y$), and multiplicative ($K(x, y) = xy$) coalescent processes, which allow an explicit analytic solution of the Smoluchowski equation; see [2, 25, 31] and references therein. Much can be learned about coalescent processes from studying the corresponding time-oriented tree graphs that describe the history of cluster merging.

This paper focuses on the trees generated by a Kingman's coalescent and its finite version, Markus-Lushnikov coalescent process, with a finite number N of particles, as N approaches infinity. We establish a particular form of the Horton self-similarity, called here *root-Horton law* for the Kingman's coalescent. We also show that the tree that describes a Kingman's coalescent is combinatorially equivalent to the level-set tree of a white noise. These findings add two important classes of processes to the realm of Horton-self-similar systems. Finally, we perform numerical experiments that suggest that the Kingman's coalescent, and hence the level-set tree of a white noise, are Horton self-similar in a regular stronger sense as well as asymptotically Tokunaga self-similar.

The paper is organized as follows. The main results are summarized in Sect. 2. The Kingman's coalescent process is introduced in Sect. 3. Section 4 introduces the concepts of Horton and Tokunaga self-similarity and the related Horton-Strahler and Tokunaga ordering schemes of tree branches. The Smoluchowski-Horton system of equations that describe the dynamics of Horton-Strahler branches in Kingman's coalescent is written out in Sect. 5. Section 6 gives the proof of the existence of root-Horton law for Kingman's coalescent, together with some related results. Section 7 establishes the equivalence between the Kingman's coalescent tree and the level-set tree of a white noise. Numerical results and the corresponding conjectures are collected in Sect. 8. A Smoluchowski-Horton system for a general coalescent process with collision kernel is written in Sect. 9. Section 10 concludes.

2 Statement of results

A trajectory of a coalescent process with a finite number of particles can be naturally described by a time-oriented tree T constructed as follows. Start with N leaves that represent the initial N particles. When two (or more) particles coalesce into a cluster, merge the corresponding leaves to form an internal vertex. Similarly, when two (or more) clusters coalesce, merge the corresponding vertices. The final coalescence would form the tree root. Each vertex within T is assigned a time of the respective coalescence. The resulting time-oriented tree represents the history of the process. It is readily seen that there is one-to-one map from the trajectories of a coalescence process with N particles onto the trees with N leaves. If multiple coalescences are not allowed, the corresponding trees are binary.

In this paper we prove that the time-oriented tree generated by a Kingman's coalescent process is Horton self-similar. We will often rely on the fact that the Kingman's coalescent is a limit, in a sense to be defined, of a Markus-Lushnikov coalescent process with N particles and constant collision kernel as $N \to \infty$ (see Sect. 3). We notice also that Horton and Tokunaga analyses only consider the combinatorial structure (branching) of a tree and ignore its metric properties (in our case – the time marks). Specifically, consider the asymptotic proportion

$$\mathcal{N}_k = \lim_{N\to\infty} N_k/N$$

of the number N_k of branches of Horton-Strahler order k in the Markus-Lushnikov coalescent of size N with constant collision kernel. Our construction of infinite tree in Sect. 7 allows one to interpret $\mathcal{N}_k$ also as the proportion of branches of order k in the infinite tree that corresponds to the Kingman's coalescent.

We show in Sect. 5 that

$$\mathcal{N}_k = \frac{1}{2}\int_0^\infty g_k^2(x)\,dx,$$

where the sequence $g_k(x)$ solves the following system of ordinary differential equations:

$$g'_{k+1}(x) - \frac{g_k^2(x)}{2} + g_k(x)g_{k+1}(x) = 0, \quad x \geq 0$$

with $g_1(x) = 2/(x+2)$, $g_k(0) = 0$ for $k \geq 2$. We also establish the following equivalent relation:

$$\mathcal{N}_k = \int_0^1 \left[1-(1-x)\,h_{k-1}(x)\right]^2 dx,$$

where

$$h'_{k+1}(x) + h_k^2(x) - 2h_k(x)h_{k+1}(x) = 0, \quad x \in [0,1]$$

with $h_0 \equiv 0$, $h_1 \equiv 1$, and $h_k(0) = 1$ for $k \geq 1$. The following theorem is proven in Section 6.

Theorem 1. *The tree associated with a Markus-Lushnikov coalescent with N particles and constant collision kernel is asymptotically Horton self-similar as $N \to \infty$. Namely,* $\lim\limits_{k\to\infty} (\mathcal{N}_k)^{\frac{1}{k}} = R$, *with* $1/4 \leq R \leq 1/2$.

In Section 7 we establish the following equivalence that expands the above Horton self-similarity to the level-set tree representation of time series of i.i.d. random variables, i.e. the white noise.

Theorem 2. *Level-set tree of a discrete white noise of length N is combinatorially equivalent to the tree generated by a Markus-Lushnikov coalescent process on N particles with constant collision kernel.*

Construction of an infinite tree for Kingman's coalescent as a limit of finite trees for Markus-Lushnikov process of increasing size (see Sect. 7) leads to the following result.

Corollary. *The infinite tree generated by the Kingman's coalescent is Horton self-similar, in the sense defined in Theorem 1.*

Finally, our numerical experiments in Sect. 8 suggest that the tree corresponding to a Kingman's coalescent obey a stronger version of Horton as well as asymptotic Tokunaga self-similarity. In addition we estimate the Horton exponent R and Tokunaga parameters (a, c). The following conjecture summarizes the results of our numerical experiments.

Conjecture. *The tree associated with Kingman's coalescent process is Horton self-similar with*

$$\lim_{k\to\infty} \frac{\mathcal{N}_{k+1}}{\mathcal{N}_k} = \lim_{k\to\infty} (\mathcal{N}_k)^{\frac{1}{k}} = R \quad \text{and} \quad \lim_{k\to\infty} \frac{\mathcal{N}_k}{R^k} = const.,$$

where $R = 0.328533\ldots$ *and Tokunaga self-similar, asymptotically in k:*

$$\lim_{i\to\infty} T_{i,i+k} =: T_k \quad \text{and} \quad \lim_{k\to\infty} \frac{T_k}{c^{k-1}} = a$$

for some positive a and c.

3 Kingman's coalecent

We start by considering a general *finite coalescent process defined by collision kernel* [31], also called *Marcus-Lushnikov process* [2, 20, 19]. The process begins with N particles (clusters) of mass one. The cluster formation is governed by a collision rate kernel $K(i, j) = K(j, i) > 0$, $1 \leq i, j \leq N - 1$. Namely, a pair of clusters with masses i and j coalesces at the rate $K(i, j)/N$, independently of the other pairs, to form a new cluster of mass $i + j$. The process continues until there is a single cluster of mass N.

Formally, for a given N consider the space of non-negative partitions

$$\Pi_N = \{(n_1, \ldots, n_N) \text{ such that } \sum_{i=1}^{N} n_i = N\}.$$

The Markus-Lushnikov process $\mathbf{ML}^{(N)}(t)$ with a positive symmetric collision kernel $K(i,j)$ is a continuous-time finite-state Markov chain on partitions Π_N. The process only can make transitions of the form

$$(n_1,\dots,n_N) \to (n_1,\dots,n_i-1,\dots,n_j-1,\dots,n_{i+j}+1,\dots,n_N)$$

at homogeneous transition rates $K(i,j)\,n_i\,n_j/N$. Here n_k represents the number of clusters of mass k in the Marcus-Lushnikov process.

The above construction uses different state space for each N. To define coalescence on the same state space, which is useful for studying the limit behavior as $N\to\infty$, we introduce infinite partitions

$$\mathcal{X} = \{(x_1 \geq x_2 \cdots \geq 0) \text{ such that } \sum_{i=1}^{\infty} x_i = 1\}.$$

A *stochastic coalescent* $\mathbf{X}(t) = (X_i(t), i \geq 1)$ is, informally, a continuous-time Markov chain with transitions

$$\begin{gathered}(x_1,\dots,x_{i-1},x_i,x_{i+1},\dots,x_{j-1},x_j,x_{j+1},\dots) \to \\ (x_1,\dots,x_i+x_j,\dots,x_{i-1},x_{i+1},\dots,x_{j-1},x_{j+1},\dots)\end{gathered}$$

that happen at homogeneous rates $K(x_i,x_j)$. We refer to [2, 31] for a formal definition. In case of a homogeneous kernel

$$K(cx,cy) = c^{\gamma}\,K(x,y), \quad 0 < c,x,y < \infty, \gamma \geq 0,$$

we have [2]

$$X_i(t) = N^{-1}\,\mathbf{ML}_i^{(N)}(t\,N^{1-\gamma}),$$

where $\mathbf{ML}_i^{(N)}(t)$ denotes the mass of the ith largest cluster in the Markus-Lushnikov process.

The *Kingman's* coalescent $\mathbf{K}(t)$ can be defined [2] as a unique version of the stochastic coalescent with kernel $K(i,j)=1$ and the initial condition

$$\lim_{t\downarrow 0}(\text{a.s.})X_1(t) = 0.$$

The Kingman's coalescent $\mathbf{K}(t)$ is a unique limit of the Markus-Lushnikov process as $N\to\infty$. Formally [2], if

$$N^{-1}\,\mathbf{ML}_1^{(N)}(0) \xrightarrow{d} 0, \quad \text{as } N \to \infty$$

then

$$\left\{N^{-1}\,\mathbf{ML}^{(N)}(Nt), 0<t<\infty\right\} \xrightarrow{d} \{\mathbf{K}(t), 0<t<\infty\}.$$

The history of a coalescent process can be naturally described by a binary tree with N leaves and $N-1$ internal vertices, each of which corresponds to an individual collision. We study here the self-similarity of this tree as $N\to\infty$.

Let $X_k(t)$ be the mean number of clusters of mass k per unit volume at time $t \geq 0$ in a coalescent process with collision kernel $K(i,j)$. Then, at time $t \geq 0$, the average number of pairs of clusters coalescing into a cluster of mass k per unit time per unit volume is

$$\frac{1}{2}\sum_{j=1}^{k-1} K(j,k-j)X_j(t)X_{k-j}(t),$$

and the average number of clusters of mass k coalescing with clusters of mass $\neq k$ per unit time per unit volume is

$$X_k(t)\sum_{j:\ j\neq k} K(k,j)X_j(t).$$

Finally, the average number of clusters of mass k coalescing with clusters of the same mass k is

$$\lim_{N\to\infty}\frac{1}{N}\left[2\binom{N\,X_k(t)}{2}\frac{K(k,k)}{N}\right] = X_k^2(t)K(k,k).$$

The *Smoluchowski coagulation equations* describe evolution of $X_k(t)$ for $k \geq 1$ as $N \to \infty$ via a system of ODEs

$$\frac{d}{dt}X_k(t) = \frac{1}{2}\sum_{j=1}^{k-1} K(j,k-j)X_j(t)X_{k-j}(t) - X_k(t)\sum_{j=1}^{\infty} K(k,j)X_j(t) \tag{1}$$

with initial conditions $X_k(0) = \delta_1(k)$, using Kronecker $\delta_1(k)$; see [2] and references therein. The Smoluchowski ODEs account for the number of clusters of masses j and $k-j$ that coalesce to new clusters of mass k, and for the clusters of mass k that coalesce with other clusters to form clusters of masses exceeding k. We will employ a similar approach in the case of Smoluchowski-Horton ODEs for Kingman's coalescent in Section 5.

4 Self-similar trees

This section describes the essential types of self-similarity for trees based on the Horton-Strahler and Tokunaga schemes for ordering nodes in a rooted binary tree. Historically, the presented approach was introduced by Horton [11] for ordering hierarchically organized river tributaries; the methods was later refined by Strahler [39] and further expanded by Tokunaga [41] to include so-called side-branching (see Sect. 4.2 below).

4.1 Planar rooted trees

A *graph* $\mathcal{G} = (V,E)$ is a collection of vertices (nodes) $V = \{v_i\}$, $1 \leq i \leq N_V$ and edges (links) $E = \{e_k\}$, $1 \leq k \leq N_E$. In a *simple* graph each edge is defined as an unordered pair of distinct vertices: $\forall\, 1 \leq k \leq N_E, \exists!\, 1 \leq i,j \leq N_V, i \neq j$ such that $e_k = (v_i, v_j)$ and we say that the edge k *connects* vertices v_i and v_j. Furthermore, each pair of vertices in a simple graph may have at most one connecting edge. A *tree* is a connected simple graph $T = (V,E)$ without cycles. In a *rooted* tree, one node is designated as a root; this imposes a natural *direction* of edges as well as the parent-child relationship between the vertices. Specifically, of the two connected vertices the one closest to the root is called *parent*, and the other – *child*. We consider planar (or labeled) trees, meaning that we will distinguish the children of the same parent. The total number of distinct trees with n leaves, according to the Cayley's formula, is n^{n-2}. The total number of distinct binary trees with n leaves is given by the $(n-1)$-th Catalan number [31]

$$C_{n-1} = \frac{1}{n}\begin{pmatrix} 2n-2 \\ n-1 \end{pmatrix}.$$

4.2 Horton-Strahler and Tokunaga order schemes

The Horton-Strahler ordering of the vertices of a finite rooted labeled binary tree is performed in a hierarchical fashion, from leaves to the root [24, 4, 11, 39]: (i) each leaf has order $r(\text{leaf}) = 1$; (ii) when both children, c_1, c_2, of a parent vertex p have the same order r, the vertex p is assigned order $r(p) = r + 1$; (iii) when two children of vertex p have different orders, the vertex p is assigned the higher order of the two. Figure 1(a) illustrates this definition. Formally,

$$r(p) = \begin{cases} r(c_1) + 1 & \text{if } r(c_1) = r(c_2), \\ \max\{r(c_1), r(c_2)\} & \text{if } r(c_1) \neq r(c_2). \end{cases} \tag{2}$$

A *branch* is defined as a union of connected vertices with the same order. The branch vertex nearest to the root is called the *initial vertex*, the vertex farthest from the root is called the *terminal vertex*. The order $\Omega(T)$ of a finite tree T is the order of its root, or, equivalently, the maximal order of its branches (or vertices). Let N_r denote the total number of branches of order r. An equivalent, and intuitively more appealing, definition of the Horton-Strahler orders can be given using the operation of *pruning*; see [4, 29, 49].

The Tokunaga indexing [24, 41, 29] extends upon the Horton-Strahler orders; it is illustrated in Fig. 1b. This indexing focuses on *side-branching*, which is the merging between branches of different order. Let $\tau_{ij}^{(k)}$, $1 \leq k \leq N_j$, $1 \leq i < j \leq \Omega$ denotes the number of branches of order i that join the non-terminal vertices of the k-th branch of order j. Then $N_{ij} = \sum_k \tau_{ij}^{(k)}$, $i < j$, is the total number of such branches in a tree T. The Tokunaga index $T_{ij} = N_{ij}/N_j$ is the average number of branches of order $i < j$ per branch of order j in a finite tree of order $\Omega \geq j$.

In a probabilistic setting, one considers a space of finite binary trees with a suitable probability measure. Then, N_i, N_{ij}, and $\tau_{ij}^{(k)}$ become random variables. We notice that if, for a given $\{ij\}$, the side-branch counts $\tau_{ij}^{(k)}$ are independent identically distributed random variables, $\tau_{ij}^{(k)} \overset{d}{=} \tau_{ij}$, then, by the law of large numbers,

$$T_{ij} \xrightarrow{\text{a.s.}} \mathsf{E}(\tau_{ij}) \quad \text{as } N_j \xrightarrow{\text{a.s.}} \infty,$$

where the *almost sure* convergence $X_r \xrightarrow{\text{a.s.}} \mu$ is understood as $\mathsf{P}\left(\lim_{r\to\infty} X_r = \mu\right) = 1$.

For consistency, we denote the total number of order-i branches that merge with other order-i branches by N_{ii} and notice that in a binary tree $N_{ii} = 2\,N_{i+1}$. This allows us to formally introduce the additional Tokunaga indices: $T_{ii} = N_{ii}/N_{i+1} \equiv 2$. The set $\{T_{ij}\}$, $1 \leq i \leq \Omega - 1, 1 \leq j \leq \Omega$, $i \leq j$ of Tokunaga indices provides a complete statistical description of the branching structure of a finite tree of order Ω.

Next, we define several types of tree self-similarity based on the Horton-Strahler and Tokunaga indexing schemes.

4.3 Horton and Tokunaga self-similarity laws

Let $\mathcal{N}_k$ define the *asymptotic ratio* of the branches of order k in a finite rooted tree T to the number N of the tree leaves defined as an almost sure limit when $N \to \infty$:

$$\mathcal{N}_k = \lim_{N\to\infty} \frac{N_k}{N}.$$

Horton self-similarity implies that the sequence $\mathcal{N}_k$ decreases in a regular geometric fashion with k. Formally, we define three types of Horton self-similarity.

Definition 1. A finite rooted tree T is said to obey a *Horton law* if and only if at least one of the following limits exists and is finite and positive:

$$\begin{aligned} &\text{(a) root law :} \quad \lim_{k\to\infty} \left(\mathcal{N}_k\right)^{-\frac{1}{k}} = R > 0, \\ &\text{(b) ratio law :} \quad \lim_{k\to\infty} \frac{\mathcal{N}_k}{\mathcal{N}_{k+1}} = R > 0, \\ &\text{(c) geometric law :} \quad \lim_{k\to\infty} \frac{\mathcal{N}_k}{R^k} = N_0 > 0. \end{aligned}$$

The constant R is called the *Horton exponent*.

It is easily seen that the three versions of the Horton law are consecutively more stricter: (c) $\Rightarrow$ (b) $\Rightarrow$ (a). The ratio and geometric Horton laws are usually considered in the literature (e.g., [29, 43, 11, 44, 6, 22, 49]). For instance, a well-known result is that a tree corresponding to a critical binary Galton-Watson process, also known as Shreve's random topology model, obey the geometric Horton law with $R = 4$; see [36, 14, 47, 29, 5, 4, 49].

A property that is more restrictive than a Horton law is conventionally required though to call a tree self-similar.

Definition 2. In a deterministic setting, a tree T of order Ω is called *self-similar* if its side-branching structure (i) is the same for all branches of a given order j:

$$\tau_{ij}^{(k)} =: \tau_{ij}, \quad 1 \leq k \leq N_j,\ 1 \leq i < j \leq \Omega,$$

and (ii) is invariant with respect to the branch order:

$$\tau_{i(i+k)} \equiv T_{i(i+k)} =: T_k \quad \text{for } 2 \leq i + k \leq \Omega. \tag{3}$$

Definition 3. A self-similar tree is called *Tokunaga self-similar* if it obeys an additional constraint first considered by Tokunaga [41]:

$$T_{k+1}/T_k = c \quad \Leftrightarrow \quad T_k = a\, c^{k-1} \quad a, c > 0,\ 1 \le k \le \Omega - 1. \tag{4}$$

For a tree satisfying the ratio-Horton and Tokunaga laws, one has [41, 29]:

$$R = \frac{2 + c + a + \sqrt{(2 + c + a)^2 - 8c}}{2}. \tag{5}$$

McConnell and Gupta [22] have shown that the ratio Horton law holds in a self-similar Tokunaga tree. Zaliapin [48] has shown, moreover, that geometric Horton laws hold in a Tokunaga tree and, at the same time, even weak Horton laws may not hold in a general, non-Tokunaga, self-similar tree.

In a random setting, we say that a tree T of order Ω is self-similar if $\mathsf{E}\left[\tau^{(j)}_{i(i+k)}\right] =: T_k$ for $1 \le j \le N_{i+k}$, $2 \le i + k \le \Omega$; and it is Tokunaga self-similar if, furthermore, the condition (4) holds.

5 Smoluchowski-Horton ODEs for Kingman's coalescent

Consider a Marcus-Lushnikov coalescent process with a constant kernel starting with N particles. We will write Smoluchowski-type ODEs for the Horton-Strahler branches of the tree that describes the process and consider asymptotic version of this equation as $N \to \infty$. We call the resulting system the *Smoluchowski-Horton* equations.

5.1 Main equation

Let $\eta(t)$ be the mean number of clusters (of any mass) at time $t \ge 0$ per unit volume. Thus $\eta(0) = N/N = 1$ and $\eta(t)$ decreases with each coalescence of clusters, at the rate of

$$\lim_{N \to \infty} \frac{1}{N^2} \binom{N\,\eta(t)}{2} = \frac{\eta^2(t)}{2}$$

per unit volume, as $1/N$ is the coalescence rate for any pair of clusters regardless of their masses. Thus $\eta(t)$ satisfies the following ODE

$$\frac{d}{dt}\eta(t) = -\frac{\eta^2(t)}{2}. \tag{6}$$

For any $j \in \mathbb{N}$ we define $\eta_j(t)$ to be the mean number of clusters of Horton-Strahler order j per unit volume at time t. Initially, each particle represents a leaf of Horton-Strahler order 1. Thus, the initial conditions are set to be, using Kronecker's delta notation

$$\eta_j(0) = \delta_1(j).$$

Recall that according to the definition of Horton-Strahler order, connecting two branches of orders i and j, where $i \neq j$, would result in a branch of order $\max\{i, j\}$. Connecting two branches of the same Horton-Strahler order, $i = j$, would result in a branch of order $j + 1$. The evolution of Horton-Strahler orders is hence very different from the evolution of cluster sizes. We describe this evolution using ODEs as follows.

Observe that at any time t, the average number of pairs of clusters of Horton-Strahler order $j - 1$ that coalesce into clusters of order j, per unit time per unit volume is

$$\lim_{N\to\infty} \frac{1}{N^2} \binom{N\,\eta_{j-1}(t)}{2} = \frac{\eta_{j-1}^2(t)}{2}.$$

Thus $\frac{\eta_{j-1}^2(t)}{2}$ is the instantaneous rate of increase of $\eta_j(t)$.

The number of clusters of order j will decrease by one if a cluster of order j will coalesce with a cluster of order strictly higher than j, and will decrease by two if a cluster of order j will coalesce with another cluster of order j. The average number of clusters of order j coalescing with clusters of order strictly greater than j per unit time per unit volume is

$$\eta_j(t)\,\left(\eta(t) - \sum_{k=1}^{j} \eta_k(t)\right),$$

and the average number of pairs of coalescing clusters of order j per unit time per unit volume is

$$\lim_{N\to\infty} \frac{1}{N^2} \binom{N\,\eta_j(t)}{2} = \frac{\eta_j^2(t)}{2}.$$

Thus the instantaneous rate of decay of $\eta_j(t)$ is

$$\eta_j(t)\,\left(\eta(t) - \sum_{k=1}^{j} \eta_k(t)\right) + \eta_j^2(t).$$

Now we can write the rates-in and the rates-out for the clusters of Horton-Strahler order via the following Smoluchowski-Horton system of ODEs:

$$\frac{d}{dt}\eta_j(t) = \frac{\eta_{j-1}(t)}{2} - \eta_j(t)\,\left(\eta(t) - \sum_{k:k<j} \eta_k(t)\right) \tag{7}$$

with the initial conditions $\eta_j(0) = \delta_1(j)$ stated earlier. Here we let $\eta_0 \equiv 0$.

We showed that $\eta_j(t)$ has the instantaneous rate of increase $\frac{\eta_{j-1}^2(t)}{2}$. The total per volume number of clusters of Horton-Strahler order j is therefore

$$\mathcal{N}_j = \delta_1(j) + \int_0^\infty \frac{\eta_{j-1}^2(t)}{2} dt.$$

In a tree generated by the Kingman's coalescent process, the above quantity $\mathcal{N}_j$ represents the asymptotic ratio for the number of branches of order j relative to the total number of leafs N.

It is not hard to compute the first few terms of sequence $\mathcal{N}_k$ by solving equations (6) and (7) in the first three iterations:

$$\mathcal{N}_1 = 1, \quad \mathcal{N}_2 = \frac{1}{3}, \quad \text{and} \quad \mathcal{N}_3 = \frac{e^4}{128} - \frac{e^2}{8} + \frac{233}{384} = 0.109686868100941\dots$$

Hence, we have $\mathcal{N}_2/\mathcal{N}_1 = 1/3$ and $\mathcal{N}_3/\mathcal{N}_2 = 3 \times \left(\frac{e^4}{128} - \frac{e^2}{8} + \frac{233}{384}\right) = 0.3290606043028\dots$ Our numerical results in Sect. 8 yield, moreover,

$$\lim_{k\to\infty} (\mathcal{N}_k)^{\frac{1}{k}} = \lim_{k\to\infty} \frac{\mathcal{N}_k}{\mathcal{N}_{k-1}} = 0.328533\dots$$

In the next section we prove the existence of the asymptotic limit $\lim\limits_{k\to\infty} (\mathcal{N}_k)^{\frac{1}{k}}$ for the root-Horton law.

6 Existence of the root-Horton law and related results

Let $g_1(t) = \eta(t)$ and $g_j(t) = \eta(t) - \sum\limits_{k:\ k<j} \eta_k(t)$ be the asymptotic number of clusters of Horton order j or higher at time t. We can rewrite (7) via g_j using $\eta_j(t) = g_j(t) - g_{j+1}(t)$ as follows

$$\frac{d}{dt} g_j(t) - \frac{d}{dt} g_{j+1}(t) = \frac{g_{j-1}(t) - g_j(t)}{2} - (g_j(t) - g_{j+1}(t))g_j(t)$$

Observe that $g_1(t) \geq g_2(t) \geq g_3(t) \geq \dots$ We now rearrange the terms, obtaining for all $j \geq 2$,

$$\frac{d}{dt} g_{j+1}(t) - \frac{g_j^2(t)}{2} + g_j(t)g_{j+1}(t) = \frac{d}{dt} g_j(t) - \frac{g_{j-1}^2(t)}{2} + g_{j-1}(t)g_j(t). \tag{8}$$

One can readily check that $\frac{d}{dt} g_2(t) - \frac{g_1^2(t)}{2} + g_1(t)\, g_2(t) = 0$; the above equations hence simplify as follows

$$g'_{j+1}(x) - \frac{g_j^2(x)}{2} + g_j(x)g_{j+1}(x) = 0 \tag{9}$$
$$\text{with} \quad g_1(x) = \frac{2}{x+2}, \quad \text{and } g_j(0) = 0 \text{ for } j \geq 2.$$

Notice that the above system of ODEs (9) is the quasilinearised Riccati equation $g'(x) = -\frac{g^2(x)}{2}$ with the initial value $g(0) = 0$ that has only a trivial solution.

Next, returning to the asymptotic ratios of the number of order-j branches to N, we observe that (8) implies that

$$\mathcal{N}_j = \int\limits_0^\infty \frac{\eta_{j-1}^2(t)}{2} dt = \int\limits_0^\infty \frac{(g_{j-1}(t) - g_j(t))^2}{2} dt = \int\limits_0^\infty \frac{g_j^2(t)}{2} dt$$

since
$$\frac{(g_{j-1}(t)-g_j(t))^2}{2}=\frac{d}{dt}g_j(t)+\frac{g_j^2(t)}{2},$$
where $\int\limits_0^\infty \frac{d}{dt}g_j(t)dt=g_j(\infty)-g(0)=0$ for $j\geq 2$. Let n_k represent the number of order-k branches relative to the number of order-$(k-1)$ branches:
$$n_k:=\frac{\mathcal{N}_k}{\mathcal{N}_{k-1}}=\frac{\int\limits_0^\infty \frac{g_k^2(x)}{2}dx}{\int\limits_0^\infty \frac{g_{k-1}^2(x)}{2}dx}=\frac{\|g_k\|^2_{L^2[0,\infty)}}{\|g_{k-1}\|^2_{L^2[0,\infty)}}.$$

Consider the following limits that represent respectively the root and the ratio asymptotic Horton laws:
$$\lim_{k\to\infty}\left(\mathcal{N}_k\right)^{\frac{1}{k}}=\lim_{k\to\infty}\left(\prod_{j=1}^k n_j\right)^{\frac{1}{k}}\qquad\text{and}\qquad \lim_{j\to\infty}n_k=\lim_{k\to\infty}\frac{\|g_k\|^2_{L^2[0,\infty)}}{\|g_{k-1}\|^2_{L^2[0,\infty)}}.$$

We expect both the limits to exist and to be equal to $0.328533\ldots$ according to our numerical results. We now establish some basic facts about g_j and n_j.

Proposition 1. *Let $g_j(x)$ be the solutions to the systems of ODEs (9). Then*

(a) $\int\limits_0^\infty \frac{g_j^2(x)}{2}dx=\int\limits_0^\infty g_j(x)g_{j+1}(x)dx,$

(b) $\int\limits_0^\infty g_{j+1}^2(x)dx=\int\limits_0^\infty (g_j(x)-g_{j+1}(x))^2dx,$

(c) $\lim\limits_{x\to\infty}xg_j(x)=2,$

(d) $n_j=\frac{\|g_j\|^2_{L^2[0,\infty)}}{\|g_{j-1}\|^2_{L^2[0,\infty)}}\leq\frac{1}{2},$

(e) $n_j=\frac{\|g_j\|^2_{L^2[0,\infty)}}{\|g_{j-1}\|^2_{L^2[0,\infty)}}\geq\frac{1}{4}.$

Proof. Part (a) follows from integrating (9), and part (b) follows from part (a). Part (c) is done by induction, using the L'Hôpital's rule as follows. It is obvious that $\lim\limits_{x\to\infty}xg_1(x)=2$. Next, suppose $\lim\limits_{x\to\infty}xg_j(x)=2$. Then
$$\lim_{x\to\infty}xg_{j+1}(x)=\lim_{x\to\infty}\frac{g_{j+1}(x)}{x^{-1}}=\lim_{x\to\infty}\frac{\frac{g_j^2(x)}{2}-g_j(x)g_{j+1}(x)}{-x^{-2}}=2\lim_{x\to\infty}xg_{j+1}(x)-2$$

implying $\lim\limits_{x\to\infty} x g_{j+1}(x) = 2$. The statement in part (d) follows from the tree construction process. A proof of part (d) using differential equations is given in the following subsection. Part (e) follows from part (a) together with Hölder inequality

$$\frac{1}{2}\|g_j\|^2_{L^2[0,\infty)} = \int\limits_0^\infty g_j(x)g_{j+1}(x)dx \le \|g_j\|_{L^2[0,\infty)} \cdot \|g_{j+1}\|_{L^2[0,\infty)},$$

which implies $\frac{\|g_{j+1}\|_{L^2[0,\infty)}}{\|g_j\|_{L^2[0,\infty)}} \ge \frac{1}{2}$.

□

6.1 Rescaling to $[0,1]$ interval

Let

$$h_k(x) = (1-x)^{-1} - (1-x)^{-2} g_{k+1}\left(\frac{2x}{1-x}\right)$$

for $x \in [0,1]$. Then $h_0 \equiv 0$, $h_1 \equiv 1$, and the system of ODEs (9) rewrites as

$$h'_{k+1}(x) = 2h_k(x)h_{k+1}(x) - h_k^2(x) \tag{10}$$

with the initial conditions $h_k(0) = 1$.

The above system of ODEs (10) is the quasilinearised Riccati equation $h'(x) = h^2(x)$ over $[0,1)$, with the initial value $h(0) = 1$. Its solution is $h(x) = \frac{1}{1-x}$. Here

$$h_k(x) \to h(x) = \frac{1}{1-x} \quad \text{and} \quad n_k = \frac{\left\|1 - h_{k+1}/h\right\|^2_{L^2[0,1]}}{\left\|1 - h_k/h\right\|^2_{L^2[0,1]}}.$$

Observe that $h_2(x) = (1+e^{2x})/2$, but for $k \ge 3$ finding a closed form expression becomes increasingly hard. Given $h_k(x)$, Eq. (10) is a linear first-order ODE in $h_{k+1}(x)$; its solution is given by $h_{k+1}(x) = \mathcal{H}h_k(x)$ with

$$\mathcal{H}f(x) = \left[1 - \int_0^x f^2(y) e^{-2\int\limits_0^y f(s)ds} dy\right] \cdot e^{2\int\limits_0^x f(s)ds}.$$

Hence, the problem we are dealing with concerns the asymptotic behavior of an iterated non-linear functional.

Using the setting of (10), we give an ODE proof to Proposition 1(d). To do so, we first need to prove the following lemma.

Lemma 1.

$$\left\|1 - h_{k+1}/h\right\|_{L^2[0,1]} = \left\|h_{k+1}/h - h_k/h\right\|_{L^2[0,1]}$$

Proof. Observe that

$$h'_{k+1}(x) + (h_{k+1}(x) - h_k(x))^2 = h_{k+1}^2(x).$$

We now use integration by parts to obtain

$$\int_0^1 \frac{(h_{k+1}(x) - h_k(x))^2}{h^2(x)} dx = \int_0^1 \frac{h_{k+1}^2(x)}{h^2(x)} dx - \int_0^1 \frac{h'_{k+1}(x)}{h^2(x)} dx$$

$$= \int_0^1 \frac{h_{k+1}^2(x)}{h^2(x)} dx - \left[\frac{h'_{k+1}(1)}{h^2(1)} - \frac{h'_{k+1}(0)}{h^2(0)} + 2\int_0^1 \frac{h_{k+1}(x)}{h(x)} dx \right]$$

$$= \int_0^1 \frac{h_{k+1}^2(x)}{h^2(x)} dx + 1 - 2\int_0^1 \frac{h_{k+1}(x)}{h(x)} dx$$

$$= \int_0^1 \frac{(1 - h_{k+1}(x))^2}{h^2(x)} dx$$

since $1/h(x) = 1 - x$. □

Alternative proof of Proposition 1(d). Notice that $h \geq \cdots \geq h_{k+1} \geq h_k \geq \cdots \geq h_0 \equiv 0$, which follows from $g_1(t) \geq g_2(t) \geq g_3(t) \geq \ldots$ The Lemma 1 implies

$$\big\|1 - h_{k+1}/h\big\|^2_{L^2[0,1]} = \big\|h_{k+1}/h - h_k/h\big\|^2_{L^2[0,1]} = \int_0^1 \left[(1 - h_k/h) - (1 - h_{k+1}/h)\right]^2 dx$$

$$= \big\|1 - h_{k+1}/h\big\|^2_{L^2[0,1]} + \big\|1 - h_k/h\big\|^2_{L^2[0,1]} - 2\int_0^1 (1 - h_k/h)(1 - h_{k+1}/h) dx$$

and therefore

$$\big\|1 - h_k/h\big\|^2_{L^2[0,1]} = 2\int_0^1 (1 - h_k/h)(1 - h_{k+1}/h) dx$$

$$= 2\big\|1 - h_{k+1}/h\big\|^2_{L^2[0,1]} + 2\int_0^1 (h_{k+1}/h - h_k/h)(1 - h_{k+1}/h) dx \geq 2\big\|1 - h_{k+1}/h\big\|^2_{L^2[0,1]}$$

yielding $\frac{1}{2} \geq \frac{\|1 - h_{k+1}/h\|^2_{L^2[0,1]}}{\|1 - h_k/h\|^2_{L^2[0,1]}} = n_{j+1}$ as in Proposition 1(d). □

It is also true that one can improve Proposition 1(d) to become a strict inequality since one can check that in fact the inequalities on h_k are strict:

$$h > \cdots > h_{k+1} > h_k > \cdots > h_0 \equiv 0.$$

6.2 Proof of the existence of the root-Horton limit

Here we present the proof of our main Theorem 1. It is based on the following two lemmas, Lemma 2 and Lemma 3, that will be proven in the following two subsections.

Lemma 2. *If the limit* $\lim\limits_{k\to\infty} \frac{h_{k+1}(1)}{h_k(1)}$ *exists, then* $\lim\limits_{k\to\infty} (\mathcal{N}_k)^{\frac{1}{k}} = \lim\limits_{k\to\infty} \left(\prod\limits_{j=1}^{k} n_j\right)^{\frac{1}{k}}$ *also exists, and*

$$\lim_{k\to\infty} (\mathcal{N}_k)^{\frac{1}{k}} = \lim_{k\to+\infty} \left(\frac{1}{h_k(1)}\right)^{\frac{1}{k}} = \lim_{k\to\infty} \frac{h_k(1)}{h_{k+1}(1)}.$$

Lemma 3. *The limit* $\lim\limits_{k\to\infty} \frac{h_{k+1}(1)}{h_k(1)} \ge 1$ *exists, and is finite.*

Theorem 1. *The limit* $\lim\limits_{k\to\infty} (\mathcal{N}_k)^{\frac{1}{k}} = \lim\limits_{k\to\infty} \left(\prod\limits_{j=1}^{k} n_j\right)^{\frac{1}{k}}$ *exists and is equal to* $\lim\limits_{k\to\infty} \frac{h_k(1)}{h_{k+1}(1)}$.

Proof. The existence and finiteness of $\lim\limits_{k\to\infty} \frac{h_{k+1}(1)}{h_k(1)}$ shown in Lemma 3 is the precondition for Lemma 2 that in turn implies the existence and finiteness of the limit $\lim\limits_{k\to\infty} (\mathcal{N}_k)^{\frac{1}{k}}$ as needed for the root-Horton law. □

6.3 Proof of Lemma 2 and related results

Proposition 2.

$$\left\|1 - h_{k+1}(x)/h(x)\right\|^2_{L^2[0,1]} \le \frac{1}{h_{k+1}(1)} \le \left\|1 - h_k(x)/h(x)\right\|^2_{L^2[0,1]}.$$

Proof. Integrating from 0 to 1 both sides of the equation

$$\frac{\frac{d}{dx}h_{k+1}(x)}{h^2_{k+1}(x)} = 1 - \frac{(h_{k+1}(x) - h_k(x))^2}{h^2_{k+1}(x)}$$

we obtain $\frac{1}{h_{k+1}(1)} = \int\limits_0^1 \frac{(h_{k+1}(x)-h_k(x))^2}{h^2_{k+1}(x)}dx$ as $h_{k+1}(0) = 1$.

Hence,

$$\frac{1}{h_{k+1}(1)} = \int\limits_0^1 \frac{(h_{k+1}(x) - h_k(x))^2}{h^2_{k+1}(x)}dx \ge \int\limits_0^1 \frac{(h_{k+1}(x) - h_k(x))^2}{h^2(x)}dx = \int\limits_0^1 \left(1 - \frac{h_{k+1}(x)}{h(x)}\right)^2 dx$$

by Proposition 1, proving the first inequality.

Now,

$$\frac{1}{h_{k+1}(1)} = \left\|1 - h_k(x)/h_{k+1}(x)\right\|^2_{L^2[0,1]} \le \left\|1 - h_k(x)/h(x)\right\|^2_{L^2[0,1]}$$

thus completing the proof. □

Proof of Lemma 2. If the limit $\lim\limits_{k\to\infty} \frac{h_{k+1}(1)}{h_k(1)}$ exists and is finite, then $\lim\limits_{k\to\infty} \left(\frac{1}{h_k(1)}\right)^{\frac{1}{k}}$ must also exist and be finite. Hence the existence and finiteness of

$$\lim_{k\to\infty} (\mathcal{N}_k)^{\frac{1}{k}} = \lim_{k\to\infty} \left(\int_0^1 \left(1 - \frac{h_k(x)}{h(x)}\right)^2 dx\right)^{\frac{1}{k}}$$

follows from Proposition 2. □

6.4 Proof of Lemma 3 and related results

In this subsection we use the approach developed by Drmota [8] to prove the existence and finiteness of $\lim\limits_{k\to\infty} \frac{h_{k+1}(1)}{h_k(1)} \geq 1$. As we observed earlier this result is needed to prove the existence and positivity of $\lim\limits_{k\to\infty} (\mathcal{N}_k)^{\frac{1}{k}} = \lim\limits_{k\to\infty} \left(\prod\limits_{j=1}^{k} n_j\right)^{\frac{1}{k}}$, the root-Horton law.

Definition 4. Given $\gamma \in (0,1]$. Let

$$V_{k,\gamma}(x) = \begin{cases} \frac{1}{1-x} & \text{for } 0 \leq x \leq 1-\gamma, \\ \gamma^{-1} h_k\left(\frac{x-(1-\gamma)}{\gamma}\right) & \text{for } 1-\gamma \leq x \leq 1. \end{cases}$$

Note that sequences of functions $h_k(x)$ and $V_{k,\gamma}(x)$ can be extended beyond $x = 1$. Here are some observations we make about the above defined functions.

Observation 1. $V_{k,\gamma}(x)$ are positive continuous functions satisfying

$$V'_{k+1,\gamma}(x) = 2V_{k+1,\gamma}(x)V_{k,\gamma}(x) - V^2_{k,\gamma}(x)$$

for all $x \in [0,1] \setminus (1-\gamma)$, with initial conditions $V_{k,\gamma}(0) = 1$.

Observation 2. Let $\gamma_k = \frac{h_k(1)}{h_{k+1}(1)}$. Then

$$V_{k,\gamma_k}(1) = h_{k+1}(1) \tag{11}$$

and

$$V_{k,\gamma}(1) = \gamma^{-1} h_k(1) \geq h_{k+1}(1) \quad \text{whenever } \gamma \leq \gamma_k. \tag{12}$$

Observation 3.

$$V_{k,\gamma}(x) \leq V_{k+1,\gamma}(x)$$

for all $x \in [0,1]$ since $h_k(x) \leq h_{k+1}(x)$.

Observation 4. Since $h_1(x) \equiv 1$ and $\gamma_1 = \frac{h_1(1)}{h_2(1)}$,

$$h_2(x) \leq V_{1,\gamma_1}(x) = \begin{cases} \frac{1}{1-x} & \text{for } 0 \leq x \leq 1-\gamma_1, \\ \gamma_1^{-1} = h_2(1) & \text{for } 1-\gamma_1 \leq x \leq 1. \end{cases}$$

The above observation generalizes as follows.

Proposition 3.

$$h_{k+1}(x) \le V_{k,\gamma_k}(x) = \begin{cases} \frac{1}{1-x} & \text{for } 0 \le x \le 1-\gamma_k, \\ \gamma_k^{-1} h_k\left(\frac{x-(1-\gamma_k)}{\gamma_k}\right) & \text{for } 1-\gamma_k \le x \le 1. \end{cases}$$

In order to prove Proposition 3 we will need the following lemma.

Lemma 4. *Functions $V_{k,\gamma}(x)$ and $h_{k+1}(x)$ have at most one positive intersection for $x \ge \gamma$ and $k \ge 1$.*

Proof. This is a proof by induction with base at $k = 1$. We need to show that if $V_{k,\gamma}(x)$ and $h_{k+1}(x)$ have at most one positive intersection, then so do $V_{k+1,\gamma}(x)$ and $h_{k+2}(x)$. Since both sequences of functions satisfy the same ODE relation (see Observation 1), we have

$$\frac{d}{dx}\left[(V_{k+1,\gamma}(x) - h_{k+2}(x)) \cdot e^{-2\int\limits_{1-\gamma}^{x} h_{k+1}(y)dy}\right]$$

$$= (2V_{k+1,\gamma}(x) - V_{k,\gamma}(x) - h_{k+1}(x)) \cdot (V_{k,\gamma}(x) - h_{k+1}(x)) \cdot e^{-2\int\limits_{1-\gamma}^{x} h_{k+1}(y)dy},$$

where $h_{k+1}(x) \le V_{k+1,\gamma}(x)$ by definition of $V_{k+1,\gamma}(x)$, and $V_{k,\gamma}(x) \le V_{k+1,\gamma}(x)$ as in Observation 3.

Now, let

$$I(x) := \int\limits_{1-\gamma}^{x} (2V_{k+1,\gamma}(s) - V_{k,\gamma}(s) - h_{k+1}(s)) \cdot (V_{k,\gamma}(s) - h_{k+1}(s)) \cdot e^{-2\int\limits_{1-\gamma}^{s} h_{k+1}(y)dy} ds.$$

Then

$$(V_{k+1,\gamma}(x) - h_{k+2}(x)) \cdot e^{-2\int\limits_{1-\gamma}^{x} h_{k+1}(y)dy} = V_{k+1,\gamma}(1-\gamma) - h_{k+2}(1-\gamma) + I(x).$$

The function $2V_{k+1,\gamma}(x) - V_{k,\gamma}(x) - h_{k+1}(x)$ is positive, and since $V_{k,\gamma}(x)$ and $h_{k+1}(x)$ have at most one positive intersection, then $I(x)$ should change the sign from positive to negative at most once as x grows from $1-\gamma$ to 1. Hence

$$V_{k+1,\gamma}(x) - h_{k+2}(x) = (V_{k+1,\gamma}(1-\gamma) - h_{k+2}(1-\gamma) + I(x)) \cdot e^{2\int\limits_{1-\gamma}^{x} h_{k+1}(y)dy}$$

should change the sign from positive to negative at most once as

$$V_{k+1,\gamma}(1-\gamma) = h(1-\gamma) > h_{k+2}(1-\gamma).$$

□

Proof of Proposition 3. Lemma 4 implies that $h_{k+1}(x)$ and $V_{k,\gamma_k}(x)$ should have no more than one positive intersection within the interval $[1-\gamma_k, 1]$. Hence, this unique positive zero is located at $x = 1$, where $h_{k+1}(1) = V_{k,\gamma_k}(1)$. Therefore, $V_{k,\gamma_k}(1-\gamma_k) > h_{k+1}(1-\gamma_k)$ implies $h_{k+1}(x) \leq V_{k,\gamma_k}(x)$ as in the statement of the proposition. ☐

Now we are ready to prove the monotonicity result.

Lemma 5.

$$\gamma_k \leq \gamma_{k+1} \qquad \textit{for all } k \in \mathbb{N}.$$

Proof. We prove it by contradiction. Suppose $\gamma_k \geq \gamma_{k+1}$ for some $k \in \mathbb{N}$. Then

$$V_{k,\gamma_k}(x) \leq V_{k,\gamma_{k+1}}(x) = \begin{cases} \frac{1}{1-x} & \text{for } 0 \leq x \leq 1-\gamma_{k+1}, \\ \gamma_{k+1}^{-1} h_k\left(\frac{x-(1-\gamma_{k+1})}{\gamma_{k+1}}\right) & \text{for } 1-\gamma_{k+1} \leq x \leq 1 \end{cases}$$

and therefore

$$h_{k+1}(x) \leq V_{k,\gamma_k}(x) \leq V_{k,\gamma_{k+1}}(x) \leq V_{k+1,\gamma_{k+1}}(x)$$

as $h_{k+1}(x) \leq V_{k,\gamma_k}(x)$ by Proposition 3.

Recall that for $x \in [1-\gamma_{k+1}, 1]$,

$$V'_{k+1,\gamma_{k+1}}(x) = 2V_{k,\gamma_{k+1}}(x)V_{k+1,\gamma_{k+1}}(x) - V^2_{k,\gamma_{k+1}},$$

where at $1-\gamma_{k+1}$ we consider only the right-hand derivative. Thus for $x \in [1-\gamma_{k+1}, 1]$,

$$\frac{d}{dx}\Big(V_{k+1,\gamma_{k+1}}(x) - h_{k+2}(x)\Big) = A(x) + B(x)\Big(V_{k+1,\gamma_{k+1}}(x) - h_{k+2}(x)\Big),$$

where $A(x) = 2V_{k+1,\gamma_{k+1}}(x) - V_{k,\gamma_{k+1}}(x) - h_{k+1}(x) \geq 0$, $B(x) = 2h_{k+1}(x) > 0$, and $V_{k+1,\gamma_{k+1}}(1-\gamma_{k+1}) - h_{k+2}(1-\gamma_{k+1}) = h(1-\gamma_{k+1}) - h_{k+2}(1-\gamma_{k+1}) > 0$. Hence

$$V_{k+1,\gamma_{k+1}}(1) - h_{k+2}(1) \geq V_{k+1,\gamma_{k+1}}(1-\gamma_{k+1}) - h_{k+2}(1-\gamma_{k+1}) > 0$$

arriving to a contradiction since $V_{k+1,\gamma_{k+1}}(1) = h_{k+2}(1)$. ☐

Corollary. *Limit* $\lim\limits_{k\to\infty} \gamma_k$ *exists.*

Proof. Lemma 5 implies γ_k is a monotone increasing sequence, bounded by 1. ☐

Proof of Lemma 3. Lemma 3 follows immediately from an observation that $\frac{h_{k+1}(1)}{h_k(1)} = \frac{1}{\gamma_k}$. ☐

7 Relation to the tree representation of white noise

This section establishes equivalence between the Kingman's coalescent and the level-set tree of a white noise. We consider separately the finite case (Markus-Lushnikov process) and infinite case (Kingman's coalescent).

7.1 Finite case

We start with recalling basic facts about tree representation of time series. For details and further results see [49]. Consider a finite time series X_k with discrete time index $k = 0, 1, \dots, N$. Let X_t (also denoted $X(t)$) be time series with continuous time $t \in [0, N]$ obtained from X_k by linear interpolation of its values. The level set $\mathcal{L}_\alpha(X_t)$ is defined as the pre-image of the function values above α:

$$\mathcal{L}_\alpha(X_t) = \{t : X_t \geq \alpha\}.$$

The level set $\mathcal{L}_\alpha$ for each α is a union of non-overlapping intervals; we write $|\mathcal{L}_\alpha|$ for their number. Notice that (i) $|\mathcal{L}_\alpha| = |\mathcal{L}_\beta|$ as soon as the interval $[\alpha, \beta]$ does not contain a value of local minima of X_t, (ii) $|\mathcal{L}_\alpha| \geq |\mathcal{L}_\beta|$ for any $\alpha > \beta$, and (iii) $0 \leq |\mathcal{L}_\alpha| \leq n$, where n is the number of the local maxima of X_t.

The *level set tree* $\textsc{level}(X_t)$ describes the topology of the level sets $\mathcal{L}_\alpha$ as a function of threshold α, as illustrated in Fig. 2. Namely, there are bijections between (i) the leaves of $\textsc{level}(X_t)$ and the local maxima of X_t, (ii) the internal (parental) vertices of $\textsc{level}(X_t)$ and the local minima of X_t (excluding possible local minima at the boundary points), and (iii) the edges of $\textsc{level}(X_t)$ and the first positive excursions of $X(t) - X(t_i)$ to right and left of each local minima t_i. The leftmost and rightmost edges of $\textsc{level}(X_t)$ may correspond to meanders, that is to a positive segments of $X(t) - X(t_i)$, rather than to excursions. It is readily seen that any function X_t with distinct values of the local minima corresponds to a binary tree $\textsc{level}(X_t)$. Please see [49] for a discussion of some subtleties related to this construction as well as for further references.

Theorem 2. *The level-set tree of a white noise of length $N < \infty$ with discrete time and continuous values is combinatorially equivalent to the tree generated by a Markus-Lushnikov coalescent process with kernel $K(i, j) = 1$ on N particles.*

Recall that the *combinatorial* equivalence refers here to considering only branching structure of a tree and ignoring the time marks.

Proof. The proof uses the duality between coalescence and fragmentation processes [2]. Specifically, a coalescence process on N particles with time-dependent collision kernel $K_t(x, y)$, $1 \leq x, y < N$ is equivalent, upon time reversal, to a discrete-mass fragmentation process of initial mass N with some *splitting kernel* $S_t(m, x)$, $1 < m \leq N$, $1 \leq x < N$. Recall that a fragmentation process starts with a single cluster of mass N. Each existing cluster of mass m splits into two clusters of masses $m - x$ and x at the splitting rate $S_t(m, x)$. See Aldous [2] for further details and the relationship between the dual collision and splitting kernels in general case.

The Marcus-Lushnikov coalescence with time-independent kernel $K(x, y) = 1$ is dual to the fragmentation process with splitting kernel [2, Table 3]

$$S_t(m, x) = \frac{2}{t\,(t + 2)}.$$

This kernel is constant with respect to the cluster mass, which means that the splitting of mass m is uniform on the interval $[1, ..., m - 1]$. The time dependence of the kernel does not affect the combinatorial structure of the fragmentation tree (and can be removed by a deterministic time change.)

The level-set tree $\textsc{level}(X_t)$ also can be considered as a tree for a fragmentation process with the initial mass N equal to the length of the time series. Indeed, by construction, each subtree of $\textsc{level}(X_t)$ with n leaves corresponds to an excursion (or meander, if we treat one of the boundaries) of length n. This subtree (as well as the corresponding excursion or meander) splits into two by the minimum of X_t at the corresponding interval. For a white noise, the location of the minimum is uniform within the interval, so is the corresponding splitting kernel. Hence, the level set tree $\textsc{level}(X_t)$ has the same combinatorial structure as the tree for a Markus-Lushnikov coalescent with constant kernel. $\square$

Remark 1. We notice that the dual splitting kernels for multiplicative and additive coalescences [2, Table 3] only differ by their time dependence, and are equivalent as functions of mass. Hence, the combinatorial structure of the respective trees is the same.

7.2 Infinite case

To treat infinite case, we recall the geometric representation of the Kingman's coalescent [2, Construction 2]. Start with placing a unit mass at each integer $i \in \mathbb{Z}$; these points represent initial particles. Clusters are formed by appearance of links of the form $(i, i+1)$ that connect initial masses to consecutively larger clusters. Each link appears at a specified time-dependent space-independent rate, independently of the other links; the particular form of time-dependence is not important for our analysis. We notice that this process is combinatorially (not considering time, only the order of link appearance) equivalent to the level-set cluster formation in a white noise. Specifically, we associate each link $(i, i+1)$ with the ith value of the white noise. The order of link appearances is now governed by the respective value of the white noise: higher values appear earlier than lower ones. Hence, the Kingman's coalescence process within each finite interval (in its geometric interpretation) can be described by a tree that is combinatorially equivalent to a tree of a finite white noise.

An infinite tree T_∞ that represents infinite white noise can be constructed as follows. Choose a local maximum of the white noise $W(t)$. The interval I_1between two consecutive local minima adjacent to the chosen local maximum correspond to a leaf of T_∞ (according to the level-set tree construction described above). We denote this leaf by T_1. Now extend the interval I_1 to the interval $I_2 \supset I_1$ between two local minima of $W(t)$ and such that it includes exactly two local maxima of $W(t)$. This extension can be done in exactly two ways: by adding the local maximum to the right or to the left of the initial one. The direction of extension does not affect the final result. The interval I_2 corresponds to the tree T_2 obtained from T_1 by adding the second leaf. Continue the extension procedure by adding another adjacent local maximum: it will correspond to interval I_3 and the tree T_3 obtained from T_2 by adding one leaf. In general, each interval I_n of the white noise that contains n consecutive local maxima corresponds to a unique branch T_n of T_∞ with n leaves. If we extend this interval to I_{n+1} by adding values to the right (of left) to include $n+1$ local maxima, then $I_n \subset I_{n+1}$ and the tree T_{n+1} that correspond to I_{n+1} is obtained from T_n by adding a single rightmost (leftmost) leaf. The precise position of this leaf within the rightmost (leftmost) branch of T_n is determined by the values of local minima within I_{n+1}.

One can also construct T_∞ using the so-called *forest of trees attached to a floor line* [31]. Namely, we start at any local minimum of the white noise and divide the trajectory into infinite sequence of finite vertically shifted positive excursions separated by intervals of monotone decrease, as discussed in [49, Section 5]. This procedure is illustrated in Fig. 3. Each excursion corresponds to a finite tree. The infinite tree T_∞ is constructed by connecting these finite trees together as illustrated in Fig. 3. The resulting tree is the same as the one constructed above by adding individual leaves.

We can summarize this discussion in the following theorem.

Theorem 3. *The Kingman's coalescent is combinatorially equivalent to the level-set tree of infinite white noise with discrete time and continuous values.*

Now we can extend the definitions of branch statistics relevant to the Horton and Tokunaga analyzes to an infinite Kingman's tree. Let us fix an arbitrary local minima of a white noise and count the other local maxima to the right and left of the chosen one using positive and negative integers $k \neq 0$, respectively. Let $N_{r,k}$, $k \in \mathbb{Z}, k \neq 0$, $r \geq 1$ denotes the number of branches of Horton-Strahler order r in the tree that corresponds to the interval of the white noise between two local minima and such that it includes all local maxima between $-k$ and k, and only those.

We notice that

$$\mathcal{N}_r = \lim_{k\to\infty} (a.s.) \frac{N_{r,k}}{2k},$$

which provides an intuitive interpretation for the quantities $\mathcal{N}_r$ in terms of an infinite tree that corresponds to the Kingman's coalescent. Clearly, our construction is another way to obtain the Kingman's coalescent as the limit of the Marcus-Lushnikov process with N particles as $N \to \infty$, in terms of the corresponding trees.

8 Numerical results

This section illustrates the theoretical results of the paper and provides further insight into self-similarity of Kingman's coalescent in a series of numerical experiments.

8.1 Horton self-similarity

Figure 4 shows (by shaded circles) the asymptotic proportion $\mathcal{N}_k$ of branches of order k in the Kingman's coalescent according to the equation

$$\mathcal{N}_k = \int_0^1 \left(1 - \frac{h_{k-1}(x)}{h(x)}\right)^2 dx. \tag{13}$$

The integral is evaluated using the numeric solution $\hat{h}_k(x)$ to the system (10). The computations are done in Matlab, using the nominal absolute tolerance of $\epsilon = 10^{-16}$ on an irregular time grid with steps decreasing towards 1 and being as small as $\Delta = 10^{-10}$. The values of the estimated $\mathcal{N}_k$ as well as the consecutive ratios $\mathcal{N}_{k+1}/\mathcal{N}_k$ are reported in Table 1 with 7 digit precision. The values of $\mathcal{N}_k$ are exponentially decreasing in such a way that the ratio

$\mathcal{N}_{k+1}/\mathcal{N}_k$ quickly converges to $R = 0.328533\ldots$. This ratio corresponds to the geometric decay

$$\mathcal{N}_k = (0.328533\ldots)^{k-1} = 10^{(k-1)\log_{10}(0.328533\ldots)} = 10^{(-0.4834\ldots)\,(k-1)}.$$

A black line in the figure illustrates geometric decay with this rate and an arbitrary offset.

The observed convergence of ratios is stronger than the root-convergence proven in our Theorem 1. We believe that in fact the strongest geometric convergence (see Definition 1) is also holds for the sequence $\mathcal{N}_k$.

Figure 4 and Table 1 report as well the number W_k of branches of order k observed in a level-set tree of a synthetic white-noise of length $N = 2^{18} = 262,144$ with Horton-Strahler order $\Omega = 11$. The values are reported with 3 significant digits. The agreement between the statistics obtained in asymptotic Kingman's coalescent and white-noise time series is very good, in accordance with our equivalence Theorem 2. We notice that the agreement is good event though we only considered a single white-noise tree; this hints at a rapid convergence of the Horton statistics as the tree size increases.

Table 1: Statistics of order-k branches for Kingman's coalescence (columns 2 and 3) and white noise (column 4).

k	$\mathcal{N}_k$	$\mathcal{N}_{k+1}/\mathcal{N}_k$	W_k/W_1
1	1.0000000	0.3333333	1.000
2	0.3333333	0.3290606	0.330
3	0.1096869	0.3285928	0.109
4	0.0360423	0.3285402	0.0356
5	0.0118413	0.3285344	0.0116
6	0.0038903	0.3285338	0.00392
7	0.0012781	0.3285337	0.00134
8	0.0004200	0.3285337	0.00043
9	0.0001380	0.3285337	0.00014
10	0.0000453	0.3285337	0.0000458
11	0.0000149	-	0.0000114

8.2 Tokunaga self-similarity

To write out the equations for the Tokunaga indexes T_{ij}, we observe that the asymptotic ratio of N_{ij} to N for $i < j$ is given by the original ODE (7) as

$$\mathcal{N}_{ij} = \int\limits_0^\infty \eta_i(t)\eta_j(t)dt = \int\limits_0^\infty \left(g_i(x) - g_{i+1}(x)\right)\left(g_j(x) - g_{j+1}(x)\right)\,dx.$$

This can be rewritten using the rescaled equations in (10) as

$$\mathcal{N}_{i-1,j-1} = 2\int_0^1 \frac{(h_i(x) - h_{i+1}(x))(h_j(x) - h_{j+1}(x))}{h^2(x)}\,dx.$$

We now use (13) to obtain

$$\lim_{N\to\infty} T_{i-1,j-1} = \lim_{N\to\infty} \frac{N_{i-1,j-1}}{N_{j-1}} = \frac{\mathcal{N}_{i-1,j-1}}{\mathcal{N}_{j-1}}$$

$$= \; 2\,\frac{\int\limits_0^1 \frac{(h_i(x)-h_{i+1}(x))(h_j(x)-h_{j+1}(x))}{h^2(x)}\,dx}{\int\limits_0^1 \left(1-\frac{h_j(x)}{h(x)}\right)^2 dx}. \tag{14}$$

Table 2 reports the Tokunaga indexes evaluated numerically using (14). The integrals and functions $h_k(x)$ are evaluated in Matlab with nominal absolute tolerance $\epsilon = 10^{-20}$ and using the same grid as in computing the Horton statistics. The values are reported here with 4 digits precision. We notice that evaluation of Tokunaga indices for larger ij pairs faces numerical problems because of divergence of $h(x)$, and the associated "bursts" of $h_k(x)$, at unity.

The reported index values $T_{i,i+k}$ converge to the limit T_k as i increases. The convergence rate is very fast; the limit value, within the reported precision, is achieved for $i \geq 4$ or faster. The reported simulations suggest that the convergence rate increases with k.

Table 2: Asymptotic Tokunaga indices T_{ij} for Kingman's coalescence

	$j=2$	3	4	5	6	7	8	9
$i=1$	0.8196	0.5687	0.2641	0.0993	0.0342	0.0114	0.0038	0.0012
2		0.8234	0.5720	0.2655	0.0999	0.0344	0.0115	0.0038
3			0.8232	0.5724	0.2657	0.0999	0.0344	0.0115
4				0.8231	0.5724	0.2657	0.0999	0.0344
5					0.8231	0.5724	0.2657	0.0999
6						0.8231	0.5724	0.2657
7							0.8231	0.5724
8								0.8231

Figure 5 shows by shaded circles the "limit" Tokunaga indices T_{i9}, $i = 1,\ldots,8$, as a function of $k = 9 - i$. The figure suggests that the Tokunaga indices form a geometric series, asymptotic in k. Summing up, we conjecture that

$$\lim_{i\to\infty} T_{i,i+k} =: T_k \quad \text{and} \quad \lim_{k\to\infty} \frac{T_k}{c^{k-1}} = a$$

with $a \approx 3 \pm 0.02$, $c \approx -0.5 \pm 0.02$ obtained by fitting a geometric series to the last couple of "limit" Tokunaga indices. It must be noted that the conventional Tokunaga self-similarity assumes the geometric form of T_k starting from $k = 1$. Hence, some of the results that hold for Tokunaga self-similar trees will not hold for the asymptotically Tokunaga trees. This applies for instance to the relation (5). Interestingly, this relation is closely satisfied, with less than 0.01 relative error, for $R = 0.328533$ from the above section and $(a,c) = (0.8230, 0.6953)$ that are obtained by fitting a geometric series to the first two "limit" Tokunaga indices $T_{8,9} \approx T_1$

and $T_{7,9} \approx T_2$. This reflects the fact that the side-branching is rapidly fading off with k and the most important side-branches, that in effect form the entire tree, are those with $k \leq 2$.

We also show in Fig. 5, by black squares, the Tokunaga indices $\hat{T}_k$ estimated for the level-set trees of discrete white noise. The values $\hat{T}_k$ are obtained by averaging the empirical Tokunaga indices in 100 trees of size $N = 2^{17} = 131,072$; the typical order of such trees is $\Omega = 11$. The indices are averaged over different trees and over all pairs of subindices $\{i, i+k\}$ with $i \neq 1$. The empirical white-noise generated values match very closely the asymptotic values from Kingman's process, in accordance with our equivalence Theorem 2.

9 General coalescent processes

The ODE approach introduced in this paper can be extended to the coalescent kernels other than $K(i,j) \equiv 1$. For that we need to classify the average number of clusters of order j per unit volume, $N_j(t)$, at time t according to cluster masses. Namely, let $\eta_{j,k}(t)$ be the average number of clusters of order j and mass $k \geq 2^j$ per unit volume, at time t. Then

$$\eta_j(t) = \sum_{k=2^j}^{\infty} \eta_{j,k}(t).$$

The average number of clusters of mass k per unit volume $X_k(t) = \sum\limits_j \eta_{j,k}(t)$ at time t can be determined using the original Smoluchowski equations (1). As for the Smoluchowski-Horton ODEs, in the case of a general coalescent process with a symmetric rate kernel $K(i,j) = K(j,i)$ these ODEs can be written asymptotically as

$$\begin{aligned} \frac{d}{dt}\eta_{j,k}(t) &= \sum_{i=1}^{j-1}\sum_{\kappa=2^j}^{k-2^i} \eta_{j,\kappa}(t)\eta_{i,k-\kappa}K(\kappa, k-\kappa) \\ &+ \frac{1}{2}\sum_{\substack{k_1+k_2=k \\ k_1,k_2\geq 2^{j-1}}} \eta_{j-1,k_1}(t)\eta_{j-1,k_2}(t)K(k_1,k_2) \\ &- \eta_{j,k}(t)\sum_{\widetilde{k}=2^i}^{\infty} K(k,\widetilde{k})\left(\sum_{i=1}^{\infty}\eta_{i,\widetilde{k}}(t)\right) \end{aligned} \tag{15}$$

with the initial conditions $\eta_{1,1}(0) = 1$ and $\eta_{j,k}(0) = 0$ for all $(j,k) \neq (1,1)$.

Observe that when $K(i,j) \equiv 1$, summing the above equations (15) over index k produces the Smoluchowski-Horton ODE (7) for the average number of order-j branches $\eta_j(t)$ per unit volume in the Kingman's coalescent process.

10 Discussion

This paper establishes Horton self-similarity and states a numerical conjecture about the asymptotic Tokunaga self-similarity for a celebrated Kingman's coalescent, which is among the most studied probabilistic models of branching structures. We also establish Horton and asymptotic Tokunaga self-similarity for a tree representation of white noise. Combining the results of this study with that of Burd et al. [4] and Zaliapin and Kovchegov [49], one concludes that Horton and Tokunaga self-similarity are intrinsic properties of (i) white noise, (ii) random walk, (iii) critical binary Galton-Watson process, and (iv) Kingman's coalescent. The listed processes are believed to closely depict actual physical and biological mechanisms of diverse origin; they are hence among the essential building blocks in natural and computer sciences modeling. The results of this study and those in [4, 49] thus provide at least a partial explanation for the omnipresence of Horton and Tokunaga self-similarity in observed and modeled branching structures.

Our Theorem 1 establishes a weak, root-Horton law (see Definition 1a), while we believe that the two stronger versions (ratio and geometric) are also valid. It seems important to show rigorously at least the ratio-Horton law, because (a) the ratio-Horton law is commonly used in the applied literature, and (b) the ability to work with asymptotic ratios is necessary to tackle the Tokunaga self-similarity, which provides much stricter constraints on a branching structure.

The Smoluchowski-Horton equations (5) that form a core of the presented method and their asymptotic equivalents (9) and (10) seem to be promising for further more detailed exploration. Indeed, one may hope that the approach that refers explicitly to the Horton-Strahler orders might effectively complement conventional analysis of cluster masses. The analysis of the Smoluchowski-Horton systems can be done within the ODE framework, similarly to the present study, or within the nonlinear iterative system framework. The latter approach is still to be explored.

Finally, it is noteworthy that the analysis of multiplicative and additive coalescents according to the general Smoluchowski-Horton system (15) appears, after a certain series of transformations, to follow many of the steps implemented in this current paper for the Kingman's coalescent, with the ODE system being replaced by a suitable PDE one. These results will be published elsewhere.

References

[1] A. Ackleh, B. Fitzpatrick, T. Hallam, *Approximation and parameter estimation problems for algal aggregation models*, Math. Models Methods Appl. Sci., **4** (1984) 291–311.

[2] D.J. Aldous, *Deterministic and stochastic models for coalescence (aggregation and coagulation): a review of the mean-field theory for probabilists*, Bernoulli, **5** (1999) 3–48.

[3] Y. Ben-Zion, C. G. Sammis, *Characterization of Fault Zones.* Pure and Appl. Geophys., **160** 3-4 (2003) 677–715.

[4] G. A. Burd, E.C. Waymire, R.D. Winn, *A self-similar invariance of critical binary Galton-Watson trees,* Bernoulli, **6** (2000) 1–21.

[5] L. Devoyre, P. Kruszewski, *A note on the Horton-Strahler number for random trees*, Inform. Processing Lett., **56** (1994) 95–99.

[6] P.S. Dodds, D.H. Rothman, *Scaling, Universality, and Geomorphology*, Ann. Rev. Earth and Planet. Sci., **28** (2000) 571–610.

[7] R. Drake, *A general mathematical survey of the coagulation equation*, In G. Hidy and J. Brock (eds), *Topics in Current Aerosol Research, Part 2,* Int. Rev. Aerosol Phys. Chem. **3** 201–376, Oxford: Pergamon, (1972).

[8] M. Drmota, *The Height of Increasing Trees* Ann. Comb. **12** (2009) 373–402

[9] A. Gabrielov, W.I. Newman, D.L. Turcotte, *An exactly soluble hierarchical clustering model: inverse cascades, self-similarity, and scaling* Phys. Rev. E. **60** (1999) 5293–5300.

[10] B. Gomez, V.T. Mullen, *An experimental study of sapped drainage network development* Earth Surface Processes and Landforms, **17** (1992) 465–476.

[11] R. E. Horton, *Erosional development of streams and their drainage basins: Hydrophysical approach to quantitative morphology* Geol. Soc. Am. Bull., **56** (1945) 275–370.

[12] A. D. Howard, *Simulation of stream networks by headward growth and branching* Geographical Analysis, **3**(l) (1971) 29–50.

[13] A. Rinaldo, I. Rodriguez-Iturbe, R. Rigon *Channel Networks* Annual Rev. Earth Planet. Sci., **26** (1998) 289–327.

[14] R. Kemp, *The average number of registers needed to evaluate a binary tree optimally.* Acta Informatica, **11** (1979) 363–372.

[15] J.F.C. Kingman, *On the Genealogy of Large Populations* J. Appl. Prob., **19A** (1982) 27-43.

[16] J.F.C. Kingman, *The coalescent* Stoch. Process. Applic., **13**, 3 (1982) 235–248.

[17] J.F.C. Kingman, *Origins of the coalescent 1974–1982* Genetics, 156 (2000) 1461-1463.

[18] S. Kramer, M. Mader, *Evolution of river networks* Phys. Rev. Lett., **68**(2) (1992) 205–208.

[19] A. Lushnikov, *Coagulation in finite systems* J. Colloid and Interface Science, **65**, (1978) 276–285.

[20] A. Marcus, *Stochastic coalescence* Technometrics, **10** (1968) 133–143.

[21] J.G. Masek, D.L. Turcotte, *A Diffusion Limited Aggregation Model for the Evolution of Drainage Networks* Earth Planet. Sci. Let. **119** (1993) 379.

[22] M. McConnell, V. Gupta, *A proof of the Horton law of stream numbers for the Tokunaga model of river networks* Fractals. **16** (2008) 227–233.

[23] J. Neveu, J. Pitman, *The branching processes in a Brownian excursion* in: Lecture Notes in Mathematics, Springer-Verlag, NY, (1989), pp. 248-257.

[24] W. I. Newman, D.L. Turcotte, A.M. Gabrielov, *Fractal trees with side branching* Fractals, **5** (1997) 603–614.

[25] J.R. Norris, *Smoluchowski's coagulation equation: uniqueness, nonuniqueness and a hydrodynamic limit for the stochastic coalescent* Ann. Appl. Probab. **9**, 1 (1999), 78-109

[26] P. Ossadnik, *Branch Order and Ramification Analysis, of Large Diffusion Limited Aggregation Clusters* Phys. Rev. A., **45** (1992) 1058.

[27] M. Ossiander, E. Waymire, Q. Zhang, *Some width function asymptotics for weighted trees* Ann. Appl. Probab. **7**, 4 (1997) 972–995.

[28] J. K. Parrish, L. Edelstein-Keshet, *Complexity, Pattern, and Evolutionary Trade-Offs in Animal Aggregation* Science, **284**, No. 5411 (1999), 99–101

[29] S. D. Peckham, *New results for self-similar trees with applications to river networks* Water Resources Res. **31** (1995) 1023–1029.

[30] J. Pitman, *Coalescents with multiple collisions* Ann. Probab. **27** (1999) 1870–1902.

[31] J. Pitman, *Combinatorial Stochastic Processes* Lecture Notes in Mathematics, vol. **1875**, Springer-Verlag (2006).

[32] J. Silk, T. Takahashi, *A statistical model for the initial stellar mass function* Astrophys. J., **229** (1979) 242–256.

[33] J. Silk, S. White, *The development of structure in the expanding universe* Astrophys. J., **223** (1978) L59–L62.

[34] R. Shcherbakov, D.L. Turcotte, *Damage and self-similarity in fracture* Theor. Appl. Fract. Mech., **39**, 3 (2003) 245–258.

[35] A. E. Scheidegger, *A stochastic model for drainage patterns into an intramontane trench* International Assoc. Scientific Hydrology, **12**, 1 (1967) 15–20.

[36] R.L. Shreve, *Statistical law of stream numbers* J. Geol., **74** (1966) 17–37.

[37] R.L. Shreve, *Stream lengths and basin area in topologically random channel networks* J. Geol, **77**, (1969) 397–414.

[38] M. Smoluchowski, Drei Vorträge über Diffusion, *Brownsche Molekularbewegung und Koagulation von Kolloidteilchen* Physik. Zeit., **17**, (1916) 557–571, 585–599

[39] A. N. Strahler, *Quantitative analysis of watershed geomorphology* Trans. Am. Geophys. Un. **38** (1957) 913–920.

[40] D.G. Tarboton, R.L. Bras, I. Rodriguez-Iturbe, *The fractal nature of river networks* Water Resources Res., **24** (1988) 1317–1322.

[41] E. Tokunaga, *Consideration on the composition of drainage networks and their evolution* Geographical Rep. Tokyo Metro. Univ. **13** (1978) 1–27.

[42] D.L. Turcotte, B.D. Malamud, G. Morein, W. I. Newman, *An inverse cascade model for self-organized critical behavior* Physica, A. **268** (1999) 629–643.

[43] D.L. Turcotte, J.D. Pelletier, and W.I. Newman, *Networks with side branching in biology* J. Theor. Biol., **193** (1998) 577–592.

[44] S. Veitzer, V.K. Gupta, *Random self-similar river networks and derivations of generalized Horton laws in terms of statistical simple scaling* Water Resour. Res., **36** (2000) 1033–1048.

[45] E. Webb. *Self-similar Trees: Genesis and Statistical Properties* Honors Undergraduate Thesis, University of Nevada, Reno (2008).

[46] G. Yakovlev, W.I. Newman, D.L. Turcotte, A. Gabrielov *An inverse cascade model for self-organized complexity and natural hazards* Geophys. J. Int. **163** (2005) 433–442.

[47] I. Yekutieli and B. B. Mandelbrot, *Horton-Strahler ordering of random binary trees* J. Phys. A: Math. Gen., **27** (1994) 285

[48] I. Zaliapin, *Horton laws in self-similar trees* (2010) preprint.

[49] I. Zaliapin and Y. Kovchegov, *Tokunaga and Horton self-similarity for level set trees of Markov chains* Chaos, Solitons & Fractals, **45**, Issue 3 (2012), pp. 358–372

[50] I. Zaliapin, H. Wong, A. Gabrielov, *Inverse cascade in percolation model: Hierarchical description of time-dependent scaling* Phys. Rev. E. **71** (2006) No. 066118.

[51] I. Zaliapin, H. Wong, A. Gabrielov, *Hierarchical aggregation in percolation model* Tectonophysics **413** (2006) 93–107.

[52] R. Ziff, *Kinetics of polymerization* J. Stat. Phys., **23**, 2 (1980), 241–263.



Figure 1: Example of (a) Horton-Strahler ordering, and of (b) Tokunaga indexing. Two order-2 branches are depicted by heavy lines in both panels. The Horton-Strahler orders refer, interchangeably, to the tree nodes or to their parent links. The Tokunaga indices refer to entire branches, and not to individual vertices.



Figure 2: Function X_t (panel a) with a finite number of local extrema and its level-set tree LEVEL(X) (panel b).



Figure 3: Illustration of tree construction for an infinite time series. The time series X_t (shown on the left) is divided here into two vertically shifted excursions, marked **A** and **B** in the time axis, and one fall, depicted by the heavy segment on the time axis. The excursions correspond to the two trees represented by marked triangles on the right.



Figure 4: Horton self-similarity. Filled circles: The asymptotic ratio $\mathcal{N}_k$ of the number N_k of branches of order k to N in Kingman's coalescent, as $N \to \infty$. Black squares: The empirical ratio N_k/N_1 in a level-set tree for a single trajectory of a white noise of length $N = 2^{18}$.



Figure 5: Asymptotic Tokunaga self-similarity. Filled circles: The asymptotic Tokunaga indices T_{i9} in Kingman's coalescent, as $N \to \infty$. Black squares: The empirical Tokunaga indices averaged over 100 level-set trees for white noises of length $N = 2^{17}$.